

## INFLUENZA – At-a-Glance

<b>DEFINITION</b>	Influenza, commonly known as "the 'flu'", is an infectious disease caused by an influenza virus. Symptoms can be mild to severe. Symptoms can include pyrexia, coryzal symptoms, muscle pains, headache, sore throat, and lethargy. In the paediatric patient there may be nausea and vomiting.
<b>MANAGEMENT</b>	<p>If a flu swab is recommended, then a throat/nasopharyngeal swab needs to be collected using a viral transport medium swab.</p> <p>Isolate in a side room – use OUH 'Droplets precautions' isolation poster. Door must remain closed. If it is not possible to isolate, please seek advice from IPC team. <b>STRICT attention to Droplets infection prevention and control precautions.</b></p> <ul style="list-style-type: none"> <li>• Hand hygiene</li> <li>• Apron and gloves to be worn if in contact with blood or bodily fluids</li> <li>• HCWs assessing or caring for patients with a suspected (clinically diagnosed) or confirmed influenza are advised to wear a fluid repellent surgical mask (single use) when in close contact with the patient (within two metres)</li> <li>• FFP3 masks (single use) are required for patients undergoing aerosol-generating procedures</li> <li>• Eye protection should be worn when there is risk of exposure to blood and/or bodily fluids or during AGPs.</li> <li>• If the patient is transferred, please inform receiving staff of patient's infection status, including radiology, theatres, or other hospital/health care settings. The patient should be asked to wear a fluid repellent surgical mask if tolerated.</li> <li>• When discharging infected patient to another healthcare setting, please complete the 'Inter-Healthcare Infection Prevention and Control transfer form'.</li> </ul> <p style="text-align: center;"><b>Encourage patients to perform good respiratory hygiene and cough etiquette (Catch it, Bin it, Kill it)</b></p>
<b>CLEANING</b>	<ul style="list-style-type: none"> <li>• <b>Equipment-</b> Single use / single patient use if possible.</li> <li>• A daily enhanced clean of the side room/ bed space for a patient with influenza should be requested via the help desk and a terminal clean on discharge.</li> <li>• All frequently touched surfaces should be clutter free and cleaned three times a day.</li> <li>• All equipment used <b>MUST</b> be decontaminated using CLINELL WIPES (green) before being brought out of the patient room and after every patient use.</li> </ul>
<b>STOP ISOLATION WHEN?</b>	These precautions should be implemented for patients with suspected or confirmed influenza for 7 days after illness onset or until 24 hours after the resolution of fever and respiratory symptoms, whichever is longer, while a patient is in the hospital/health care setting. For the severely immunocompromised, discuss with Infection Prevention & Control or Microbiology/Infectious Diseases doctor.